

Phenylketonuria and BH4 Deficiencies

Nenad Blau, in collaboration with Alberto B. Burlina, Barbara K. Burton, Harvey L. Levy, Nicola Longo, Anita MacDonald, Aurora Martinez, Francjan J. van Spronsen, Beat Thöny

UNI-MED, 3rd edition 2016, 112 pp, Hardcover, ISBN 978-3-8374-1526-1, Euro 39,80

ePDF-version: ISBN 978-3-8374-5526-7, Euro 39,80



In patients with phenylketonuria (PKU), blood phenylalanine concentration during childhood is the major determinant of cognitive outcome. Thanks to newborn screening and early dietary therapy, individuals with PKU no longer experience intellectual disability. Nevertheless, some do not achieve their full potential. The establishment of uniform guidelines and improved management for PKU can lead to optimal outcomes in this metabolic disorder.

Since in 1999 it has been shown that some patients with PKU respond to the administration of tetrahydrobiopterin (BH4; sapropterin dihydrochloride) by lowering blood phenylalanine concentrations, that these patients can be treated with sapropterin dihydrochloride. Enzyme substitution therapy with phenylalanine ammonia lyase (PAL) might become a new option in the near future, along with diet and sapropterin, to reduce Phe levels and improve the clinical outcome of subjects with PKU. It has been also shown that patient's genotype determines the phenotype and helps in predicting BH4 responsiveness.

In this textbook past, present, and future efforts related to PKU and BH4 deficiencies are discussed. The reviews and scientific contributions in this book provide professionals, the patients, and their families to understand PKU within a biochemical, neurological and psychological context.

**Klinische
Lehrbuchreihe**

... Kompetenz und Didaktik!

SCIENCE